

Agterberg and Dodgson Reply

D.F. Agterberg¹ and Matthew J.W. Dodgson^{2,3}

¹*Department of Physics, University of Wisconsin-Milwaukee, Milwaukee, WI 53201*

²*Theory of Condensed Matter Group, Cavendish Laboratory, Cambridge, CB3 0HE, United Kingdom*

³*Institut de Physique, Université de Neuchâtel, 2000 Neuchâtel, Switzerland*

A reply to the Comment by Mineev and Champel.

In their Comment [1], Mineev and Champel have argued that our results are incorrect because of the simultaneous neglect of both non-linear and non-local terms in our theory. We disagree with this statement and point out that while Ref. [1] argue that these terms should be included, they do not give a convincing physical argument that these terms are important at low fields. In the following, we explicitly show that non-linear and non-local terms can be safely ignored *near* H_{c1} where our results are valid. Furthermore, our results agree with the assertion of Ref. [1] that there is no true A to B phase transition in the vortex phase. In Ref. [2], we refer to the diverging correlation length that describes the A to B transition in the *Meissner* phase.

To justify our claims, we give the missing steps between Eq. 2 and Eq. 3 of Ref. [2]. The expression used for Ψ_- in the A phase was

$$(\xi_{A,-}^{-2} - \mathbf{D}^2)\Psi_-(\mathbf{r}) = \frac{\tilde{\kappa}}{\kappa}(D_x D_y + D_y D_x)\Psi_+(\mathbf{r}) \quad (1)$$

where $D_i = \partial_i + i2\pi A_i/\Phi_0$. Eq. 3 in Ref. [1] is found by setting the $-\mathbf{D}^2$ operator to zero. This operator cuts off the divergence that Ref. [1] point out in their Eq. 3. We work with $\tilde{\kappa}/\kappa \ll 1$ and within a London approach. Taking $\Psi_+(\mathbf{r}) = |\Psi_+|e^{i\phi}$, $\Psi_- = e^{i\phi}\tilde{\Psi}_-$, and defining the superfluid velocity as $\mathbf{v} = \nabla\phi + 2\pi\mathbf{A}/\Phi_0$ we get

$$(\xi_-^{-2} + \mathbf{v}^2 - 2i\mathbf{v} \cdot \nabla - \nabla^2)\tilde{\Psi}_-(\mathbf{r}) = |\Psi_+|\frac{\tilde{\kappa}}{\kappa}(i\partial_x v_y + i\partial_y v_x - 2v_x v_y). \quad (2)$$

This last equation shows that $\tilde{\Psi}_-$ will be proportional to v (for small v). Consequently, for large vortex separations, the non-linear term in v on the right hand side and the terms with v on the left hand side of Eq. 2 can be ignored ($v \propto e^{-r/\lambda}/\sqrt{r\lambda}$ and the derivatives give a factor $1/\lambda \gg v$ for large r). After Fourier transforming and using the Maxwell relation $\mathbf{v} = \frac{2\pi\lambda^2}{\Phi_0}\nabla \times \mathbf{B}$, Eq. 2 becomes

$$\tilde{\Psi}_-(\mathbf{q}) = |\Psi_+|\frac{\tilde{\kappa}}{\kappa}\frac{q_x^2 - q_y^2}{\xi_{A,-}^{-2} + \mathbf{q}^2}\frac{2\pi\lambda^2}{\Phi_0}B(\mathbf{q}). \quad (3)$$

Using this expression in the free energy and minimizing with respect to $B(\mathbf{q})$ gives the novel London equation that forms the basis of our results

$$\left[1 + \lambda^2 \mathbf{q}^2 + \frac{\lambda^2 \tilde{\kappa}^2}{\kappa^2} \frac{(q_x^2 - q_y^2)^2}{\xi_{A,-}^{-2} + \mathbf{q}^2}\right]B(\mathbf{q}) = 0. \quad (4)$$

This gives a 4-fold symmetry to the structure of a flux line out to the distance $\xi_{A,-}$, which diverges at the $A \rightarrow B$ transition temperature.

Using Eq. 3, $|\Psi_-(\mathbf{r})| \lesssim \frac{\tilde{\kappa}}{\kappa}|\Psi_+|\frac{2\pi\lambda^2 B(\mathbf{r})}{\Phi_0}$ (take $q_x^2 - q_y^2 \rightarrow \mathbf{q}^2$ in Eq. 3). This, with $\beta_1|\psi_+|^2/\kappa = 1/\xi_+^2$ and $2\pi\lambda^2 B(\mathbf{r})/\Phi_0 = \lambda^2/d^2$, we find:

$$\frac{\langle \beta_1 |\Psi_-|^4 \rangle}{\langle \kappa |\mathbf{D}\Psi_-|^2 \rangle} \lesssim \frac{\tilde{\kappa}^2}{\kappa^2} \frac{\lambda^2}{\xi_{A,+}^2} \frac{\lambda^4}{d^4} \quad (5)$$

where d is the distance between vortices. Ref. [1] uses this result as their Eq. 5. H_{c1} marks a second order phase transition between the Meissner and the vortex phase. At this transition $B = 0$, which implies that as $H \rightarrow H_{c1}$, $d \rightarrow \infty$. Consequently, sufficiently near H_{c1} , the non-linear term can be safely neglected. Mineev and Champel argue that d is cutoff when $d \approx \lambda/\sqrt{\ln(\lambda/\xi_{A,+})}$. Presumably, this accounts for the very small range of H over which B goes from H_{c1} to zero (or d goes from λ to ∞). Nevertheless, this range is experimentally accessible, and furthermore, for flat plate-like samples the demagnetization factors force the applied field $H_{app} \propto B$, thus making the large d limit

even more accessible. It is also argued that when Eq. 6 in Ref.[1] is not satisfied, then non-local corrections must be included (these are terms that are $O(\mathbf{q}^4)$ or larger in Eq. 4). To counter this, we point out that Eq. 4 implies new physics at low fields (for small q and large d) while non-local terms give new physics at high fields (for large q and small d). In fact, we find non-local terms are not important when $\tilde{\kappa}/\kappa > \frac{\xi_{A,+}}{\lambda} \frac{\lambda}{d}$. Consequently, non-local terms also become negligible in the large d limit.

DFA was supported by an award from Research Corporation. MJWD was supported by an EPSRC Advanced Fellowship AF/99/0725.

- [1] V.P. Mineev and T. Champel, preceding comment.
- [2] D.F. Agterberg and M.J.W. Dodgson, Phys. Rev. Lett. **89**, 017004 (2002).